SLAYER **MULTI-PURPOSE INSECT SPRAY**



DESCRIPTION

SLAYER is an EPA registered water base aerosol true multipurpose insect killer. SLAYER contains the synthetic pyrethroid permethrin for quick knockdown and up to four weeks residual control of crawling and flying insects. Non-staining formula controls bedbugs and many other crawling insects. Controls houseflies, mosquitoes and many other flying insects. For use on dogs and on premises to control fleas and ticks. For use on livestock to control flies, fleas and gnats. Controls garden insects on ornamentals and plants. Authorized for use in animal quarters and milk houses.

BENEFITS - FEATURES

- * Kills and Controls Bedbugs
- * Broad-Spectrum Spray
- * Kills on Contact
- * Will not harm plants
- * No fragrance
- * Up to 4 Week Residual

ADVANTAGES

- * Killing Action Is Stronger Than Natural Pyrethrins
- * Kills Flying And Crawling Insects
- * No fragrance

WHERE TO USE

Anywhere that flying and crawling insects are being a pest

RECOMMENDED FOR

- * Restaurants
- * Hotels
- * Nursing Homes
- * Office Buildings
- * Food Processing Plants
- * Schools
- * Hospitals
- * Dairy Farms

DIRECTIONS

SHAKE WELL BEFORE USING

SHAKE WELL BEFORE USING
Hold container upright and spray from a distance of 12 to 15 inches. Remove birds and cover fish aquariums before spraying. Spray until surface is just damp, but not to the point of wetness or run-off. Avoid excessive application. Keep children and/or pets out of treated areas during treatment. Do not apply this product to or near children's toys. Vacate room after treatment and ventilate before reoccupying. Cover or remove fishbowls.
HOUSEFLIES, FRUIT FLIES, MOSQUITOES, GNATS, BLACK FLIES, SKIPPER FLIES, BLOW FLIES AND SMALL FLYING MOTHS: Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Upon reentry, open windows and ventilate for at least 2 hours. Repeat

reatment once a month.

COCKROACHES, WATER BUGS, PALMETTO BUGS, SILVERFISH, FIREBRATS, CRICKETS, SCORPIONS, SPIDERS, MILLIPEDES, CENTIPEDES, SOWBUGS AND PILLBUGS: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray when ever possible. Upon reentry, open windows and ventilate for at least 2 hours. Repeat treatment once a menth.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat treatment once a month. INDIAN MEAL MOTHS, ANGOUMOIS GRAIN MOTHS:

Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Upon reentry, open windows and ventilate for at least 2 hours. Repeat treatment once a month.

EXPOSED STAGES OF CONFUSED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, RED FLOUR BEETLES, RICE WEEVILS, GRANARY WEEVILS, GRAIN MITES: Spray into cracks and crevices, processing and handling equipment, around cartons or containers or

other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Upon reentry, open windows and ventilate for at least 2 hours. Repeat treatment once a month. Infested stored food

products in the home should be disposed of.

CARPET BEETLES: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment once a

BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Allow all sprayed articles to dry thoroughly before reuse. After mattress is dry cover

with mattress cover and sheet. Do not use mattress without cover. Repeat treatment once a month.

TO CONTROL CLOTHES MOTHS AND CARPET BEETLES on wool and other proteinaceous fibers: Remove infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated by spraying lightly, keeping nozzle at least three feet from fabric to avoid staining. Allow all sprayed articles to dry thoroughly before

IN ANIMAL QUARTERS AND MILK ROOMS: For rapid control of House Flies, Mosquitoes, Gnats, and Small-Flying Moths. Remove animals prior to application. Close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed. Upon reentry, open windows and ventilate for at least 2 hours. Repeat

treatment once a month.

TO TREAT LIVESTOCK: To protect beef cattle, dairy cattle, sheep, hogs and horses from attacks of Stable Flies, Horse Flies, Deer Flies, Face Flies, House Flies, Horn Flies, Mosquitoes, and Gnats. Spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from Face Flies, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome. flies are troublesome.

FOR SWINE: Do not ship animals for slaughter within 5 days of last treatment

CONTROL OF BEDBUGS AND MITES IN POULTRY

HOUSES, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

FLEAS AND TICKS (ADULTS AND LARVAE): Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering, with special attention to cracks and crevices around baseboards, and with a person of the property and window and door frames. Infested upholstered furniture may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.

Inconspicuous piace before use.

TO CONTROL FLEAS AND TICKS ON DOGS. Starting at the tail, stroke against the lay of the hair, spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Repeat application every two weeks if necessary.

FOR THE CONTROL OF GARDEN INSECTS: For use on

Repeat application every two weeks if necessary. FOR THE CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray, as well as Whiteflies, Aphids, exposed Thrips, Spiders, Red Mites and Leafminers. May be used on plants, as well as a multipurpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas. May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, eunonymus, fuschia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias. Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lady Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Army Worms. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur. Notice: Buyer assumes all responsibility for safety and use not in accordance with directions.

SPECIFICATIONS

Spray pattern.....Fine mist Appearance.....Milky emulsion Flammability.....Nonflammable Wetting Ability......Good

Rinsability......Complete Fragrance.....None Shelf life.....1 year min. Net weight......16 oz.

ACTIVE INGREDIENTS:

*Permethrin. 0.250% 1(3-phenoxyphenyl)methyl (+/-)cis/trans 3-(2,2-dichloroethenyl)-2, 2-dimethycyclopropanecarboxylate (Cis/Trans ratio: Min. 35 % (+/-) cis and max. 65% (+/-)

Another Exclusive Product of

